

AD-A149 390

AN INFRARED STUDY OF ISOTOPIC EXCHANGE DURING
METHANATION OVER SUPPORTED. (U) AUBURN UNIV AL DEPT OF
CHEMISTRY M A HENDERSON ET AL. 07 JAN 85 TR-3
N00014-83-K-0637

1/1

UNCLASSIFIED


F/G 18/2

NL

END

FILMED

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS													
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Unlimited													
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE															
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 3		5. MONITORING ORGANIZATION REPORT NUMBER(S)													
6a. NAME OF PERFORMING ORGANIZATION Auburn University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research-Code 413													
6c. ADDRESS (City, State and ZIP Code) Department of Chemistry (Dr. S. D. Worley) Auburn University Auburn University, AL 36849		7b. ADDRESS (City, State and ZIP Code) 800 North Quincy St. Arlington, VA 22217													
8a. NAME OF FUNDING/SPONSORING ORGANIZATION ONR	8b. OFFICE SYMBOL (If applicable) Code 413	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-K-0637													
8c. ADDRESS (City, State and ZIP Code) See 7b		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO.</td><td>PROJECT NO.</td><td>TASK NO.</td><td>WORK UNIT NO.</td></tr><tr><td></td><td></td><td></td><td></td></tr></table>		PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.								
PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT NO.												
11. TITLE (Include Security Classification) An Infrared Study of Isotopic Exchange During Methanation Over Supported Rhodium Catalysts: An Inverse Spillover Effect NR634-844															
12. PERSONAL AUTHOR(S) M. A. Henderson and S. D. Worley															
13a. TYPE OF REPORT Technical	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Yr., Mo., Day) 85 Jan 7	15. PAGE COUNT 12												
16. SUPPLEMENTARY NOTATION Accepted for publication by J. Phys. Chem.															
17. COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr></table>		FIELD	GROUP	SUB. GR.										18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Infrared Spectroscopy, Isotopic Exchange, Methanation, Supported Rhodium Catalysts, Carbon Dioxide, Inverse Spillover Effect.	
FIELD	GROUP	SUB. GR.													
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Infrared spectroscopy has been employed to study the reduction of carbon dioxide on supported catalyst films. The investigation included isotopic labeling using D ₂ as the reduction gas. Isotopic exchange was observed for both CO ₂ /D ₂ and CH ₄ /D ₂ mixtures. The mechanisms of this isotopic exchange involves migration of hydrogen from the support to the Rh sites, an "inverse spillover effect". A key intermediate in the dissociation of CO ₂ on the supported Rh films was a carbonyl hydride species.															
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified													
22a. NAME OF RESPONSIBLE INDIVIDUAL S. D. Worley		22b. TELEPHONE NUMBER (Include Area Code) 205-826-4043	22c. OFFICE SYMBOL												

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

AD-A149 390

DTIC FILE COPY

85 01 08 052

H.W. T.II 1/31/85
JAN 31, 1985

TR

UNCLASSIFIED

PAGE 28

PP 00 A149390 (U)

FIELD/GROUP 000000

UNCLASSIFIED TITLE

AN INFRARED STUDY OF ISOTOPIC EXCHANGE DURING METHANATION OVER SUPPORTED RHODIUM CATALYSTS; AN INVERSE SPILLOVER EFFECT

ABSTRACT

(U) INFRARED SPECTROSCOPY HAS BEEN EMPLOYED TO STUDY THE REDUCTION OF CARBON DIOXIDE ON SUPPORTED CATALYST FILMS. THE INVESTIGATION INCLUDED ISOTOPIC LABELING USING DEUTERIUM AS THE REDUCTION GAS. ISOTOPIC EXCHANGE WAS OBSERVED FOR BOTH CARBON DIOXIDE/D2 AND METHANE/DEUTERIUM MIXTURES. THE MECHANISM OF THIS ISOTOPIC EXCHANGE INVOLVES MIGRATION OF HYDROGEN FROM THE SUPPORT TO THE RHODIUM SITES, AN INVERSE SPILLOVER EFFECT. A KEY INTERMEDIATE IN THE DISSOCIATION OF CARBON DIOXIDE ON THE SUPPORTED RH FILMS WAS A CARBONYL HYDRIDE SPECIES.

CARBON DIOXIDE

USE CARBON DIOXIDE

DISSOCIATION OF CARBON DIOXIDE

USE CARBON DIOXIDE

DISSOCIATION

INVERSE SPILLOVER EFFECT

USE INVERSION

SPILLING

METHANE DEUTERIUM MIXTURES

USE DEUTERIUM

METHANE MIXTURES

REDUCTION GAS

USE GASES

REDUCTION

SUPPORTED CATALYST FILMS

USE CATALYSTS

FILMS

CARBON DIOXIDE D2

POSTING TERMS ASSIGNED

DEUTERIUM

USE DEUTERIUM

INFRARED SPECTROSCOPY

USE INFRARED SPECTROSCOPY

ISOTOPIC EXCHANGE

USE EXCHANGE

ISOTOPES

MIGRATION OF HYDROGEN

USE HYDROGEN

MIGRATION

RHODIUM SITES

USE RHODIUM

SITES

SUPPORTED RHODIUM CATALYSTS

USE CATALYSTS

RHODIUM

PHRASES NOT FOUND DURING LEXICAL DICTIONARY MATCH PROCESS

UNCLASSIFIED

* Isotope exchange - co.
Reduction (Chemistry) - co.
Chemical dissociation - co.
Carbonyl compounds - co.
Hydrides - co.

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-83-K-0637

Task No. NR 634-844

TECHNICAL REPORT NO. 3

An Infrared Study of Isotopic Exchange During
Methanation Over Supported Rhodium Catalysts;
An Inverse Spillover Effect

by

M. A. Henderson and S.D. Worley
Department of Chemistry
Auburn University
Auburn University, AL 36849

Prepared for Publication

in

J. Phys. Chem.

August 6, 1984



Reproduction in whole or in part is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited.

An Infrared Study Of Isotopic Exchange During Methanation Over Supported Rhodium Catalysts; An Inverse Spillover Effect

M.A. Henderson and S.D. Worley*

Department of Chemistry, Auburn University, Auburn University, AL 36849

Abstract

Infrared spectroscopy has been employed to study the reduction of carbon dioxide on supported catalyst films. The investigation included isotopic labeling using D_2 as the reduction gas. Isotopic exchange was observed for both CO_2/D_2 and CH_4/D_2 mixtures. The mechanism of this isotopic exchange involves migration of hydrogen from the support to the Rh sites, an "inverse spillover effect". A key intermediate in the dissociation of CO_2 on the supported Rh films was a carbonyl hydride species.

deuterium

methane

carbon dioxide

rhodium

*Author to whom correspondence should be addressed.

i -

Introduction

A substantial amount of effort in these laboratories has been devoted recently to the study of the hydrogenation of carbon dioxide over supported rhodium catalysts. Of considerable interest and importance in this reaction is its likely first step, the dissociation of CO_2 on the supported Rh. The dissociation of CO_2 on Rh, or lack thereof, is a controversial topic at this time, with Somorjai and coworkers favoring a high dissociation probability (10^{-1}) at 300 K,¹ while Weinberg² and Goodman and coworkers³ suggest a low probability (10^{-15} and 10^{-11} at 300 K, respectively). Very recently Solymosi and Kiss⁴ have concluded that the dissociation probability at 300 K is low on clean Rh (111), but enhanced by the presence of small amounts of hydrogen gas or boron impurity. Henderson and Worley have reached a similar conclusion for supported Rh catalyst films.⁵ Part of the preliminary evidence concerning CO_2 dissociation on Rh/ TiO_2 and Rh/ Al_2O_3 obtained in these laboratories⁵ was derived from isotopic labeling studies. The purpose of this letter is to present those data in more detail and to examine the hydrogen spillover phenomena for supported Rh catalyst films using infrared spectroscopy as a probe.

Experimental


The supported Rh catalysts used in this study were prepared as thin films on CaF_2 infrared windows by spraying slurries of $\text{RhCl}_3 \cdot 3\text{H}_2\text{O}$, support material, water, and acetone onto the windows under mild heating to remove the solvents. Complete details concerning this method of sample preparation and subsequent reduction procedures have been presented previously concerning infrared studies of CO adsorbed on supported Rh⁶ and the hydrogenation of CO over supported Rh.⁷ The support materials were

alumina (Degussa Aluminum Oxide C, $100 \text{ m}^2\text{g}^{-1}$) and titania (Degussa Titanium Dioxide P25, $50 \text{ m}^2\text{g}^{-1}$). The gases employed (H_2 , D_2 , CO_2 , CH_4) were of the highest purity obtainable from Matheson; the H_2 , D_2 and CH_4 were passed through liquid nitrogen traps before use. The Pyrex infrared cell used in this investigation was similar to those employed in our earlier studies^{6,7} except that it was designed such that it could be heated to ca. 523 K in situ in a Perkin-Elmer model 983 infrared spectrometer. The spectrometer was operated at a resolution of 4.5 cm^{-1} in the 2000 cm^{-1} region, with a Perkin-Elmer infrared data station being employed to facilitate data processing and storage. All experiments were run several times with the results obtained being entirely reproducible.

Results and Discussion

Figure 1 illustrates the effects of heating a mixture of CO_2 and D_2 over a 10% by weight Rh/ TiO_2 catalyst film. The film produced by the slurry-deposition technique mentioned in the Experimental Section was heated for 12 hr at 523 K at 2×10^{-6} Torr to remove as much hydrogen (from adsorbed H_2O and OH moieties) from the support as possible. However, it is well-established that this treatment at only 523 K is not sufficient to remove all hydrogen from the support. Then the film was reduced at 470 K by four successive doses of D_2 at 70 Torr for time intervals of 5, 5, 10, and 20 min, each reduction cycle being followed by evacuation to 10^{-5} Torr. Following evacuation to 2×10^{-6} Torr at 298 K, a 66 torr 1:4 mixture of CO_2 : D_2 was introduced into the cell at 298 K, and spectrum 1a and Fig. 1 was recorded. The primary infrared bands observed in 1a (with the exception of the CO_2 gas-phase band at 2349 cm^{-1}) at 2704, 2626, and 2527 cm^{-1} can be attributed to the deuterated support (OD groups); any remaining

OH on the support was not of sufficient concentration to give observable OH infrared bands near 3600 or 1600 cm^{-1} . A weak band in spectrum 1a at ca. 2020 cm^{-1} is also evident, even at 298 K. This band can be assigned to a carbonyl deuteride species formed



as a result of CO_2 dissociation. The corresponding carbonyl hydride band at low coverage for CO_2 dissociation enhanced by H_2 appears at ca. 2030 cm^{-1} .⁵ The reproducible 10 cm^{-1} shift to lower frequency caused by the deuterium isotope effect is consistent with isotopic substitution two bonds removed from the CO moiety. It should be noted that Solymosi and coworkers first suggested the existence of such a carbonyl hydride species during CO_2 dissociation.⁸ We believe that the isotope effect observed in this study confirms the suggestion of Solymosi and coworkers, and thus negates the arguments of Iizuka and Tanaka⁹ which essentially stated that bands observed in the 2020-2040 cm^{-1} region of the infrared spectra of dissociated CO_2 over supported Rh were due to a linear CO species at low coverage rather than a carbonyl hydride.

Following heating the sample at 503 K for 1 hr, spectrum 1b was recorded. The sharp band at 2258 cm^{-1} can be assigned to a C-D stretching mode of gas-phase CD_4 ,¹⁰ the expected methanation product. A band at 996 cm^{-1} could also be resolved using data subtraction which can be attributed to CD_4 . The carbonyl deuteride band is shifted to higher frequency (2029 cm^{-1}) due to higher coverage of this species; the corresponding carbonyl hydride band for a sample treated identically, except using H_2 instead of D_2 , appears at 2039 cm^{-1} . Bands at 2778 and 2720 cm^{-1} can be assigned to

vibrational modes for gas-phase D_2O and HDO , respectively.¹⁰ Also present were small sharp bands at 2988 cm^{-1} and 2141 cm^{-1} . The 2988 cm^{-1} band is indicative of the presence of a small amount of CHD_3 . The 2141 cm^{-1} band does not grow in intensity upon further heating and is removed upon evacuation. At this time we can not propose an assignment for this band.

Spectra 1c and 1d show the further development of the bands mentioned above upon further heating at 503 K. New bands at 3010, 1234, 1089, and 1035 cm^{-1} can be assigned to vibrational modes for gas-phase CH_2D_2 . Upon evacuation at 298 K, spectrum 1e was obtained showing only the carbonyl deuteride surface species and broad bands indicative of a deuterated support. It should be noted that an analogous series of experiments for 10% Rh/ Al_2O_3 gave the CD_4 , CHD_3 , and CH_2D_2 products, but at a slower rate. Also a 0.5% Rh/ Al_2O_3 catalyst film gave qualitatively the same results, although the 0.5% sample actually produced the isotopically labeled methanes at a slightly greater rate than did the 10% sample indicating that small Rh ensembles can be important in the methanation of CO_2 .

In a similar set of experiments a 10% Rh/ TiO_2 film reduced with D_2 , heated at 523 K in the presence of 70 Torr D_2 for 21 hr, and then evacuated at 298 K to 1×10^{-6} Torr, was exposed to a 66 torr 1:4 mixture of $CO_2:H_2$ at 298 K. At 298 K the infrared spectrum revealed no measurable OD bands, yet a broad band at 1610 cm^{-1} characteristic of a hydrated (H) support appeared. Nevertheless, upon heating to 503 K for 3 hr, infrared bands at 2199 and 1156 cm^{-1} characteristic of CH_3D , as well as those (3014 and 1304 cm^{-1}) for CH_4 , were observed.

Figure 2 shows the infrared spectra for a series of experiments involving a CH_4/D_2 mixture over a 10% Rh/ TiO_2 film. For these experiments the film was heated and reduced in D_2 . Then 100 Torr of a 1:1 mixture of

CH₄:D₂ was added at 298 K, and spectrum 2a was recorded. At 298 K only the 3014 and 1304 cm⁻¹ bands for CH₄ are evident. However, after heating to 503 K for 30 min (2b) and 6 hr (2c), bands at 2986, 2258, 2199, 1234, 1156, 1089, 1034, and 994 cm⁻¹ clearly indicate the presence of CH₃D, CH₂D₂, CHD₃, and CD₄, in addition to CH₄. Evacuation at 298 K led to the removal of all of these bands (2d). Similar results were obtained for a 0.5% Rh/TiO₂ film.

All of these experiments suggest that isotopic scrambling is occurring on the supported Rh films. To discover whether this scrambling was related to the Rh, or just the support materials, the CO₂ reduction experiments and the CH₄/D₂ mixture experiments were repeated for TiO₂ and Al₂O₃ films containing no Rh. The resulting infrared spectra show no infrared bands characteristic of any of the methanes other than the CH₄ when a CH₄/D₂ mixture was used at the temperatures employed in these experiments. Utiyama and coworkers¹¹ have reported that isotopic exchange in a CH₄/D₂ mixture does not occur over TiO₂ until temperatures greater than 773 K. However, other workers have observed CH₄/D₂ exchange over Al₂O₃ at low temperature (near ambient).¹²⁻¹⁴ The sample pretreatment procedures in these experiments all involved heating to temperatures in excess of 750 K in O₂ or air before the exchange was attempted. Evidently our mild pretreatment procedure did not activate either TiO₂ or Al₂O₃ for isotopic exchange. Thus, it is apparent that all isotopic exchange in these experiments was induced by the presence of the Rh on the supports.

We believe that isotopic exchange in these experiments is a direct result of H or D migration from the support material to the Rh atoms or clusters. Otherwise, CHD₃ and CH₂D₂ could not be formed from a gaseous mixture of D₂ and CO₂ over the supported Rh films as in Fig. 1. There has

been much interest recently in hydrogen and deuterium "spillover" from metals to supports.¹⁵ There have also been suggestions of "reverse spillover" of D₂ onto Pt from TiO₂¹⁶ and "back spillover" of protons onto Pt from TiO₂,¹⁷ among other similar observations.¹⁸⁻²¹ We prefer to term the phenomenon as an "inverse spillover effect", and to our knowledge, this is the first documented example of such an effect for supported Rh catalysts. Furthermore, we believe that the "inverse spillover effect" significantly enhances the dissociation of CO₂ on supported Rh catalysts in the absence of H₂ gas.^{5,22}

Acknowledgements

We gratefully acknowledge the support of the Office of Naval Research for this work. We also thank F. Solymosi for access to reference 4 in advance of publication.

References

- (1) B.A. Sexton and G.A. Somorjai, J. Catal., **46** 167 (1977); D.G. Castner, B.A. Sexton, and G.A. Somorjai, Surf. Sci., **71**, 519(1978); D.G. Castner and G.A. Somorjai, Surf. Sci., **83**, 60(1979); L.H. Dubois and G.A. Somorjai, Surf. Sci., **88**, L13(1979); L.H. Dubois and G.A. Somorjai, Surf. Sci., **91**, 514(1980); L.H. Dubois and G.A. Somorjai, Surf. Sci., **128**, L231(1983); see also the numerous references quoted therein.
- (2) W.H. Weinberg, Surf. Sci., **128**, L224(1983).
- (3) D.W. Goodman, D.E. Peebles, and J.M. White, Surf. Sci., **140**, L239(1984).
- (4) F. Solymosi and J. Kiss, Surf. Sci., in press; Chem. Phys. Letters, in press.
- (5) M.A. Henderson and S.D. Worley, Surf. Sci., submitted.
- (6) C.A. Rice, S.D. Worley, C.W. Curtis, J.A. Guin, and A.R. Tarrer em. Phys., **74**, 6487 (1981); S.D. Worley, C.A. Rice, G.A. Mattson, C.W. Curtis, J.A. Guin, and A.R. Tarrer, J. Chem. Phys., **76**, 20 (1982); S. Worley, C.A. Rice, G.A. Mattson, C.W. Curtis, J.A. Guin, and A.R. Tarrer, Phys. Chem., **86**, 2714 (1982).
- (7) S.D. Worley, G.A. Mattson, and R. Caudill, J. Phys. Chem., **87**, 1671(1983).
- (8) F. Solymosi, A. Erdohelyi, and M. Kocsis, J. Catal. **65**, 428(1980); F. Solymosi, A. Erdohelyi, and T. Bansagi, J. Catal. **68**, 371(1981); F. Solymosi, A. Erdohelyi, and T. Bansagi, J. Chem. Soc. Faraday Trans. 1, **77**, 2645 (1981); A. Erdohelyi, M. Kocsis, T. Bansagi, and F. Solymosi, Acta. Chim. Acad. Scien. Hung. Tomus, **111** 591(1982).
- (9) T. Iizuka and Y. Tanaka, J. Catal., **70**, 449(1981); T. Iizuka, Y. Tanaka, and K. Tanabe, J. Catal. **76**, 1(1982).
- (10) G. Herzberg, "Molecular Spectra and Molecular Structure II. Infrared and Raman Spectra of Polyatomic Molecules", D. Van Nostrand, Inc., New York, 1945.
- (11) M. Utiyama, H. Hattori, and K. Tanabe, J. Catal., **53** 237(1978).
- (12) J.G. Larson and W.K. Hall, J. Phys. Chem., **69**, 3080(1965).
- (13) B.D. Flockhart, S.S. Uppal, and R.C. Pink, Trans. Faraday Soc., **67**, 513(1971).
- (14) P.J. Robertson, M.S. Scurrall, and C. Kemball, J. Chem. Soc. Faraday Trans. 1, **71**, 903(1975).

- (15) For a few examples see D.D. Beck and J.M. White, J. Phys. Chem., **88**, 175 (1984); D.D. Beck and J.M. White, J. Phys. Chem., **88**, 2764(1984); R.R. Cavanagh and J.T. Yates, J. Catal., **68**, 22(1981); J.C. Conesa and J. Soria, J. Phys. Chem., **86**, 1392(1982); T.M. Apple and C.R. Dybowski, J. Catal., **71**, 316(1981); S.J. DeCanio, T.M. Apple, and C.R. Dybowski, J. Phys. Chem. **87**, 194(1983).
- (16) D.D. Beck, A.O. Bawagan, and J.M. White, J. Phys. Chem., **88**, 2771(1984).
- (17) H. Jiehan, H. Zupei, S. Yongze, and W. Hongli, Stud. Sur. Sci. Catal., **17**, 53(1983).
- (18) P.A. Sermon and G.C. Bond, J. Chem. Soc. Faraday Trans. 1, **72**, 745(1976).
- (19) T. Fleisch and R. Abermann, J. Catal., **50**, 268(1977).
- (20) T. Huizinga and R. Prins, J. Phys. Chem., **85**, 2156(1981).
- (21) S.J. DeCanio, J.B. Miller, J.B. Michel, and C.R. Dybowski, J. Phys. Chem., **87**, 4619(1983).
- (22) A referee has suggested that an alternative explanation for our observations is a water gas shift sequence in which D₂O produced from deuteration of CO₂ exchanges with the support to produce HDO which then reacts with CO to give CO₂ and HD. The resulting HD would then react with CO₂ on Rh to produce the CHD₃ and CH₂D₂ observed. While this explanation is possible, we believe that an inverse spillover effect is more plausible because of the substantial amount of the deuterated products observed in a reactor in which D₂ would be in tremendous excess relative to any HD produced. Also, we observe infrared bands for CHD₃ before those for HDO appear.

Figure Captions

Figure 1. Infrared spectra of a 10% Rh/TiO₂ film (4.8 mg cm⁻²) following heating at 523 K for 12 hr at 2x10⁻⁶ Torr, reduction in D₂ at 470 K, evacuation at 298 K to 2x10⁻⁶ Torr, and then: (a) introduction of 66 Torr of a 1:4 mixture of CO₂:D₂ at 298 K, (b) heating at 503 K for 1 hr, (c) heating at 503 K for 3 hr, (d) heating at 503 K for 4.5 hr, (e) following evacuation at 298 K.

Figure 2. Infrared spectra of a 10% Rh/TiO₂ film (5.1 mg cm⁻²) following reduction in D₂ at 470 K and evacuation at 298 K to 2x10⁻⁶ Torr, and then: (a) introduction of 100 Torr of a 1:1 mixture of CH₄:D₂ at 298 K, (b) heating at 503 K for 30 min, (c) heating at 503 K for 6 hr, (d) following evacuation at 298 K.




Fig. 2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

ABSTRACTS DISTRIBUTION LIST, 634

Dr. K. N. Nicholas
Department of Chemistry
University of Oklahoma
Norman, Oklahoma 73019

Dr. Aaron Fletcher
Code 3852
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. J. Cooper
Code 6130
Naval Research Laboratory
Washington, D.C. 20375

Dr. M. C. Baird
Department of Chemistry
Queen's University
Kingston, Canada
K7L 3N6

Dr. N. N. Lichtin
Department of Chemistry
Boston University
685 Commonwealth Avenue
Boston, Massachusetts 02215

Dr. R. Eisenberg
Department of Chemistry
University of Rochester
Rochester, New York 14627

Dr. N. Cooper
Department of Chemistry
Harvard University
Cambridge, Massachusetts 02138

Dr. S. D. Worley
Department of Chemistry
Auburn University
Auburn, Alabama 36830

Dr. L. Vaska
Department of Chemistry
Clarkson College
Potsdam, New York 13676

Dr. R. A. Huggins
Materials Science and
Engineering Department
Stanford University
Stanford, California 94305

END

FILMED

2-85

DTIC